

Reši parcialno diferencialno enačbo  $\frac{\partial^2}{\partial x^2} u(x, y) = 0$

## Matematika 4

4. vaja

B. Jurčič Zlobec<sup>1</sup>

<sup>1</sup>Univerza v Ljubljani,  
Fakulteta za Elektrotehniko  
1000 Ljubljana, Tržaška 25, Slovenija

Matematika FE, Ljubljana, Slovenija 17. april 2013

►  $\frac{\partial^2}{\partial x^2} u(x, y) = 0.$

►  $\frac{\partial}{\partial x} u(x, y) = f(y).$

►  $u(x, y) = f(y)x + g(y).$

Reši parcialno diferencialno enačbo  $\frac{\partial^2 u(x, y)}{\partial x \partial y} = 0$

Reši parcialno diferencialno enačbo  $\frac{\partial^2 u(x, y)}{\partial x \partial y} + \frac{\partial u(x, y)}{\partial x} = 0$

►  $\frac{\partial^2}{\partial x \partial y} u(x, y) = 0.$

► Uvedemo novo spremenljivko  $u(x, y)_x = v(x, y).$

►  $v(x, y)_y + v(x, y) = 0 \rightarrow v(x, y) = f'(x)e^{-y}.$

►  $u_x(x, y) = f'(x)e^y \rightarrow u(x, y) = f(x)e^{-y} + g(y).$

►  $u(x, y) = f(x)e^{-y} + g(y).$

►  $\frac{\partial^2}{\partial x \partial y} u(x, y) + \frac{\partial}{\partial x} u(x, y) = 0.$

► Uvedemo novo spremenljivko  $u(x, y)_x = v(x, y).$

►  $v(x, y)_y + v(x, y) = 0 \rightarrow v(x, y) = f'(x)e^{-y}.$

►  $u_x(x, y) = f'(x)e^y \rightarrow u(x, y) = f(x)e^{-y} + g(y).$

►  $u(x, y) = f(x)e^{-y} + g(y).$

Reši parcialno diferencialno enačbo  $\frac{\partial^2 u(x,y)}{\partial x^2} + u(x,y) = 0$

Reši parcialno diferencialno enačbo  $\frac{\partial^2 u(x,y)}{\partial y^2} + u(x,y) = 0$

- $\frac{\partial^2}{\partial x^2} u(x,y) + u(x,y) = 0.$
- $u(x,y) = f(y) \cos x + g(y) \sin x.$
- $\frac{\partial}{\partial y} u(x,y) = x$
- $u(x,y) = xy + f(x).$

Reši parcialno diferencialno enačbo  
$$\frac{\partial^2 u(x,y)}{\partial x \partial y} + \frac{\partial u(x,y)}{\partial x} + x + y = 0$$

- $\frac{\partial^2 u}{\partial x \partial y} u(x,y) + \frac{\partial u}{\partial x} u(x,y) + x + y = 0.$
- Uvedemo novo spremenljivko  $u(x,y)_x = v(x,y) \rightarrow v(x,y)_y + v = -x - y.$
- Rešitev homogene enačbe  $v_{hy} + v_h = 0$  je  $v_h = f'(x)e^{-y}.$
- Partikularna rešitev  $\bar{v}(x,y) = C(y)e^y.$
- $\bar{v}'(x,y)_y + \bar{v}(x,y) = -x - y \rightarrow C'(y)e^{-y} - C(y)e^{-y} + C(y)e^{-y} = -x - y \rightarrow C'(y) = (-x - y)e^y \rightarrow C(y) = -xe^y - ye^y + e^y \rightarrow \bar{v}(x,y) = -x - y + 1.$
- $u(x,y)_x = v(x,y) = f'(x)e^{-y} - x - y + 1 \rightarrow$
- $u(x,y) = f(x)e^{-y} - \frac{x^2}{2} - xy - x + g(y).$

Reši parcialno diferencialno enačbo  $\frac{\partial u(x,y)}{\partial x} = 2xyu(x,y)$

- Vzamemo, da je  $y$  konstanta.
- Navadna diferencialna enačba ima ločljive spremenljivke.
- Pišemo  $v(x) = u(x,y) \rightarrow v' = 2xyv \rightarrow \frac{dv}{v} = 2xy dx.$
- $\ln|v| = x^2y + \ln|C| \rightarrow v(x) = C e^{x^2y}.$
- $u(x,y) = C(y)e^{x^2y}.$

Reši parcialno diferencialno enačbo  $\frac{\partial u(x,y)}{\partial x} + x \frac{\partial^2 u(x,y)}{\partial x^2} = y$

Reši parcialno diferencialno enačbo  $\frac{\partial^2 u(x,y)}{\partial y^2} + \frac{\partial u(x,y)}{\partial y} = xy$

- Uvedemo novo spremenljivko in vzamemo, da je y konstanta.
- $v(x) = u(x,y)_x \rightarrow v + xv' = y \rightarrow x \frac{dv}{dx} = y - v$ .
- Ločimo spremenljivke  $\frac{dv}{y-v} = \frac{dx}{x}$ .
- $\ln|y-v| = \ln|x| + \ln|C| \rightarrow y - v = Cx \rightarrow v = y - Cx$ .
- $u(x,y)_x = y + f(y)x \rightarrow u(x,y) = xy + f(y)\frac{x^2}{2} + g(y)$ .
- $u(x,y) = xy + f(y)\frac{x^2}{2} + g(y)$ .

$$\blacksquare u(x,y) = \frac{1}{2}xy^2 - xy + f(x)e^{-y} + g(x).$$

- Uvedemo novo spremenljivko in vzamemo, da je x konstanta.
- $v(y) = u(x,y)_y \rightarrow v' + v = xy$ .
- Rešitev homogene enačbe je  $v(y) = Ce^{-y}$ .
- Nastavek za partikularno rešitev  $\bar{v}(y) = C(y)e^{-y}$ .
- $C'(y)e^{-y} = xy \rightarrow C'(y) = xy e^y \rightarrow C(y) = x(ye^y - e^y)$
- $\bar{v}(y) = xy - x \rightarrow v(y) = xy - x + Ce^{-y}$ .
- $u(x,y)_x = xy - x + f(x)e^{-y} \rightarrow u(x,y) = \frac{1}{2}xy^2 - xy + f(x)e^{-y} + g(x)$ .
- $u(x,y) = \frac{1}{2}xy^2 - xy + f(x)e^{-y} + g(x)$ .

Vpelji nove spremenljivke in reši enačbo  
 $\frac{\partial^2 u(x,y)}{\partial x^2} - 2 \frac{\partial^2 u(x,y)}{\partial x \partial y} + \frac{\partial^2 u(x,y)}{\partial y^2} = 0, t = x, z = x + y$ .

Vpelji nove spremenljivke in reši enačbo  
 $x \frac{\partial u(x,y)}{\partial x} - y \frac{\partial u(x,y)}{\partial y} = 2u(x,y), t = x^2, z = xy$

- $\frac{\partial u}{\partial x} = 2x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial z} \rightarrow \frac{\partial u}{\partial y} = x \frac{\partial u}{\partial z}$ .
- $\frac{\partial u}{\partial x} = 2\sqrt{t} \frac{\partial u}{\partial x} + \frac{z}{\sqrt{t}} \frac{\partial u}{\partial z} \rightarrow \frac{\partial u}{\partial y} = \sqrt{t} \frac{\partial u}{\partial z}$ .
- $\sqrt{t}2\sqrt{t}u_t + \sqrt{t}\frac{z}{\sqrt{t}}u_z - \frac{z}{\sqrt{t}}\sqrt{t}u_z = 2u$
- $2tu_t = 2u \rightarrow \frac{du}{u} = \frac{dt}{t} \rightarrow u = tf(z)$
- $u(x,y) = x^2f(xy)$ .
- $u(x,y) = f(x+y)x + g(x+y)$ .

Vpelji nove spremenljivke in reši enačbo

$$\frac{\partial^2 u(x,y)}{\partial x^2} + \frac{\partial^2 u(x,y)}{\partial x \partial y} - 2\frac{\partial^2 u(x,y)}{\partial y^2} = 0, \quad t = x + y, \quad z = 2x - y$$

Poisci rešitev diferencialne enačbe  $\frac{\partial u(x,y)}{\partial x} + \frac{\partial u(x,y)}{\partial y} = 0$  v obliki  $u(x,y) = X(x)Y(y)$

- ▶  $u_x = u_t + 2u_z \quad u_y = u_t - u_z \rightarrow$
- ▶  $u_{xx} = (u_t + 2u_z)_t + 2(u_t + 2u_z)_z = u_{tt} + 4u_{tz} + 4u_{zz},$
- ▶  $u_{yy} = (u_t - u_z)_t - (u_t - u_z)_z = u_{tt} - 2u_{tz} + u_{zz},$
- ▶  $u_{xy} = (u_t + 2u_z)_t - (u_t + 2u_z)_z.$
- ▶  $u_{xx} - 2u_{xy} + u_{yy} = u_{tt} + 2u_{tz} + u_{zz} - 2u_{tz} - 2u_{zz} \rightarrow$   
 $u_{tt} = 0.$
- ▶  $u = f(z)t + g(z) \rightarrow u(x,y) = f(2x-y)(x+y) + g(2x-y).$
- ▶  $u(x,y) = f(2x-y)(x+y) + g(2x-y).$

Poisci rešitev diferencialne enačbe  $x^2 u_{xy} + 3y^2 u = 0$  v obliki  $u = XY$

- ▶  $x^2 X' Y' + 3y^2 XY = 0 \rightarrow$
- ▶  $x^2 \frac{X'}{X} \frac{Y'}{Y} + 3y^2 = 0 \rightarrow$
- ▶  $x^2 \frac{X'}{X} = -3y^2 \frac{Y}{Y'} = k,$
- ▶ kjer je  $k$  parameter nadvisen od  $x$  in  $y$ .
- ▶  $\frac{dX}{X} = k dx$  in  $\frac{dY}{Y'} = -y dy \rightarrow$
- ▶  $X(x) = Ae^{kx}$  in  $Y(y) = Be^{-ky} \rightarrow$
- ▶  $u(1,0) = Ce^k = 1$  in  $u(0,1) = Ce^{-\frac{k}{2}} = 2$
- ▶  $k = -\frac{1}{3} \log 4, \quad C = 4^{1/3}.$

Poisci rešitev diferencialne enačbe  $u_x + yu_y = 0$  v obliki  $u = XY$ , ki ustreza pogojema  $u(1,0) = 1$  in  $u(0,1) = 2$ .